



[Previous abstract](#) | [Graphical version](#) | [Text version](#) | [Next abstract](#)

Session D41 - Poster Session I.

POSTER session, Monday evening, March 17

Exhibit Hall D, Conv. Center

[D41.67] Surface Tamm States In The Propagation Of Electromagnetic Longitudinal Waves In Periodic Superlattices

Gerardo Vazquez-Fonseca (IFUNAM), Victor M. Ortega-Montiel (IFUNAM,FCUNAM), Marcelo del Castillo-Mussot (IFUNAM)

We model a system of alternating layers (made of metallic or highly doped semiconductor) and an interface of other conducting material at $z=0$ as a system of wells and barriers. The use of ABC's implies the continuity of μ_j and ν_{ρ} where j and ρ are the current density and electronic density, respectively, μ and ν might depend on material. We use the hydrodynamic model to find a dispersion relation for longitudinal electromagnetic modes of the superlattice. This relation is similar to that in the Kronig-Penney model. Using border conditions at $z=0$ we found an equation for the decaying parameter p . With the use of the equation for the decaying parameter and the dispersion relation of the superlattice we get to equation for the surface modes (also called Tamm states) for longitudinal electromagnetic waves.

▪ **Part D of program listing**



[Previous abstract](#) | [Graphical version](#) | [Text version](#) | [Next abstract](#)

Session D41 - Poster Session I.

POSTER session, Monday evening, March 17

Exhibit Hall D, Conv. Center

[D41.67] Surface Tamm States In The Propagation Of Electromagnetic Longitudinal Waves In Periodic Superlattices

Gerardo Vazquez-Fonseca (IFUNAM), Victor M. Ortega-Montiel (IFUNAM,FCUNAM), Marcelo del Castillo-Mussot (IFUNAM)

Log #7245

Abstract Submitted
for the MAR97 Meeting of
The American Physical Society

Sorting Category: 13.a

Surface Tamm States In The Propagation Of Electromagnetic Longitudinal Waves In Periodic Superlattices
GERARDO VAZQUEZ-FONSECA, IFUNAM, VICTOR M. ORTEGA-MONTIEL, IFUNAM, FCUNAM, MARCELO DEL CASTILLO-MUSSOT, IFUNAM — We model a system of alternating layers (made of metallic or highly doped semiconductor) and an interface of other conducting material at $z = 0$ as a system of wells and barriers. The use of ABC's implies the continuity of $\mu \mathbf{j}$ and $\nu \rho$ where \mathbf{j} and ρ are the current density and electronic density, respectively, μ and ν might depend on material. We use the hydrodynamic model to find a dispersion relation for longitudinal electromagnetic modes of the superlattice. This relation is similar to that in the Kronig-Penney model. Using border conditions at $z = 0$ we found an equation for the decaying parameter p . With the use of the equation for the decaying parameter and the dispersion relation of the superlattice we get to equation for the surface modes (also called **Tamm states**) for longitudinal electromagnetic waves. Start your abstract by replacing this line with your text.

☐ Prefer Oral Session
☒ Prefer Poster Session

Date submitted: 5 Dec 1996

Marcelo del Castillo-Mussot
mussot@ifunam.ifisicacu.unam.mx
IFUNAM
Member ID: 60007251

Electronic form version 1.2

■ Part D of program listing

u